

# BREAST CANCER IS THE NUMBER TWO KILLER DISEASE IN THE US

*Breast cancer is the most common cancer in American women, This also means there is a 7 in 8 chance she will never have the disease.*

Users can develop a comprehensive understanding of their risk, leading to better financial planning, a more in-depth understanding of their current portfolio, enhanced risk selection in future underwriting decisions, and improved hedging strategies. An estimated 2, men in the United States will be diagnosed with breast cancer. This means that it can be difficult to estimate each person's chance of survival. Breast Cancer Facts What is breast cancer? Being over 60 years old increases the risk of CKD, as does having a family history of it. About 40, women are expected to die from breast cancer in the U. They decided to look at perception of risk, and asked about women aged 35 to 49 about breast cancer risk versus heart disease risk. This includes women currently being treated and women who have finished treatment. Kidney disease Percentage of total deaths: 1. Furthermore, higher rates of obesity which is associated with diabetes can be another reason for the increasing rate of heart attack among young adults. The primary goal is generally to cure cancer or to considerably prolong life. As a result of this, waste from the blood remains in the body and may lead to other health problems. Type C viruses are fairly uncommon. More women are diagnosed with breast cancer than any other cancer, besides skin cancer. Damaged kidneys cannot filter blood as well as healthy kidneys. Vaccination against these HPV and hepatitis B viruses could prevent 1 million cancer cases each year<sup>3</sup>. About 2, new cases of invasive breast cancer are expected to be diagnosed in men in Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks, a systematic analysis for the Global Burden of Disease Study Screening programmes can be effective for select cancer types when appropriate tests are used, implemented effectively, linked to other steps in the screening process and when quality is assured. However, since , the number of women who have died of breast cancer has steadily decreased thanks to early detection and treatment improvements. She notes that many of the things advocated for in each campaignâ€”upping your physical activity , maintaining a healthy body weight, eating healthy foods, and other common-sense strategiesâ€”all work well to ward off breast cancer and keep your heart happy. Also, experts measure the survival statistics every 5 years. Suicide Deaths in 47, When a person dies by suicide, they may have lived with a mental health condition â€” such as depression , anxiety , or bipolar disorder â€” for a long time.